

Thermal Imaging Techniques in Agricultural Applications



Aryalekshmi B N, R C Biradar, Mohammed Ahamed J

Abstract: Thermal imaging innovation has transformed into an important technique in all fields as a result of its non-ruinous, non-contact method to choose thermal properties and highlights of any object/zone of interest. It is a procedure of improving the permeability of objects in a dark environment by detecting the objects infrared radiation and shaping an image based on that obtained data. This method will be useful in all fields where temperature difference helps in analyzing the area/object. The potential use of this technique in the agricultural field includes irrigation scheduling, soil properties mapping, plant disease detection, crop yield estimation, evaluating the maturity of the crop, etc. This paper briefly reviews some of the works on thermal imaging for agricultural applications.

Keywords—Thermal imaging, agriculture, irrigation scheduling, remote sensing.

I. INTRODUCTION

The farming segment is the principal patron of the Indian economy. Conventional methods for ascertaining agrarian parameters are solid; however they are tedious, requires more labour, and constrained for little size territories. Thermal remote detecting gives precise information for parameter portrayal than regular procedures since it gives ceaseless airborne scope over a wide zone at a particular time interim. Texas Instruments initially produced thermal imaging for military applications in the late 1960s. It is the method of utilizing the heat emitted by an object to create an image of it or to find it. These images can be procured utilizing compact, handheld or heat sensors that are mounted on satellite or a flying machine. The quality of radiation produced by an object relies upon its surface temperature.

Thermal imaging has got various application in agriculture beginning from soil water stress estimation, crop water stress estimation, water system planning, plant disease identification, farming plastic waste location, etc. Thermal imaging framework includes thermal camera furnished with a picture securing framework, a signal processing unit, and infrared indicators. The infrared radiation discharged by the object is absorbed by the locator and is changed over into

electrical signal which is sent to signal processing framework where the data will be changed into thermal image. Thermal imaging gadgets are of two kinds to be specific cooled and uncooled. An infrared imaging framework will be viewed as in light of the picture resolution, examine speed and thermal affectability. Figure1 describes the taxonomy of thermal imaging for agricultural applications. The purpose of this paper is to review and condense the major application of thermal imaging in agricultural filed.




Fig 1: Taxonomy of thermal imaging for agricultural applications

II. THERMAL IMAGING APPLICATION IN AGRICULTURAL SECTOR

Vegetation reflectance spectrum qualities rely upon leaves properties; however, the measure of reflected energy for a wavelength relies upon thickness and leaf pigmentation [1], [2]. Thermal imaging in the agricultural application we have mainly classified into seven sections as shown in the taxonomy (fig1) as irrigation scheduling, crop maturity and yield mapping,

Revised Manuscript Received on October 30, 2019.

* Correspondence Author

Aryalekshmi B N*, School of ECE, REVA University, Bangalore, India. Email: aryalks88@gmail.com

Dr. R C Biradar, School of ECE, REVA University, Bangalore, India. Email: dir.ece@reva.edu.in

Mohammed Ahamed J, RRSC-South, NRSC, Bangalore, India. Email: mohammedahamed_j@nrsc.gov.in

© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an open access article under the CC-BY-NC-ND license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

soil properties mapping, field tile mapping, agricultural plastic waste estimation, crop residue cover, and tillage mapping and plant disease detection. Irrigation scheduling can again be classified into soil moisture detection, crop water-stress monitoring and evapotranspiration and drought stress monitoring. Challenges and opportunities of satellite and UAV (Unmanned Aerial Vehicle) based agricultural applications have been studied in various journals and books[3]–[7]. For thermal remote sensing mostly, Landsat data is used. In [8], [9] a guide to Landsat pre-processing is given which can be used for monitoring irrigation scheduling, agricultural plastic waste estimation, etc. In this paper, we will be focusing on the works done on various agricultural applications, which is given in the next section.

III. IRRIGATION SCHEDULING USING THERMAL IMAGING

In numerous agrarian areas, restriction underway is principal because of poor water system timing and deficient utilization of water; accordingly, the water system is basic to take care of harvest water demand. As the plant goes into water pressure condition, their stomata start to close making the crop water stress to rise [10]. In [11] demonstration of the outcomes acquired with a low-cost UAV system for farming, applications are done. Soil moisture detection, crop water stress monitoring (canopy temperature), evapotranspiration (ET), and drought monitoring, soil salinity detection, and crop/fruit yield detection will come under thermal imaging irrigation scheduling application, which will be discussed in the next section.

A. Soil moisture detection

Soil moisture content is one of the vital factors in hydrological forms, which impact the trading of water and vitality transitions at the land surface and air interface [12], [13]. For numerous environmental studies, an exact gauge of spatial and transient variety of soil dampness is required. Recent innovative improvement in satellite remote sensing has exhibited that moisture content in soil can be evaluated by various remote sensing procedure with its strength as well as weakness. Remote sensing summary for estimation of moisture content in soil is given in **Table 1**. Change in soil dampness has a high impact on the thermal properties of the soil. The triangle method and soil moisture index are commonly used thermal imaging approaches for measuring soil moisture [14]. Nonstop checking of moisture content in soil is important as it gives supplements to plant development, controls soil temperature, impacts agricultural works like irrigation, planting, etc. [15]. Vegetation that is well water will be having less radiant temperature compared to vegetation subject to water stress [16]. In [17] the author shows that Perpendicular Soil Moisture Index (PSMI) utilizing data from Landsat satellite images is unequivocally connected with observed soil dampness. The exhibition of assessing the spatial appropriation of surface soil dampness in ranch/farm planted with oats by utilizing thermal images is appeared in [18]. In agro-production, the spatial inconsistency of soil dampness can be in charge of low or spatially variable trim yields, as soil dampness is required to make supplements dissolvable for plant assimilation [19]. Another way to estimate soil dampness in light of evaporative fraction (EF) recovered from optical/thermal infrared MODIS

information was displayed in [20]. The results can be possibly used to downscale soil dampness estimations got from dormant microwave remote recognizing techniques. Nevertheless, the model required further enhancements to represent extraordinary soil dampness conditions.

A comparison of thermal infrared imaging to thermocouple estimation and evaluation of environmental impact was studied in [21]. The result shows that TIR estimations can be amended to evaluate canopy temperature all the more precisely and that the significance of these correction increments under progressively outrageous surrounding conditions, for example, colder periods. [22] Contemplated LiDAR and thermal images combination for the ground-based 3D portrayal of fruit trees where the framework conveys numerous perspectives of the plantation, providing the client a 3D perspective of the warm conduct of the grove. Utilizing a managed classifier, the executed calculation characterizes focuses on the LiDAR estimations that compare to the canopy. Later thermal camera provides the thermal data by performing coordinating method between such focuses. Utilizing the Iterative Closest point, the whole plantation or just a segment of the grove is remade.

Methods for utilizing thermal infrared radiation to survey spatial variations in soil water accessibility likewise have utility in precision agriculture [23]. The connection between the soil salinity and canopy temperature, standardized distinction vegetation index and enhanced vegetation index was done in [24]. A remote sensing based trim water pressure index for irrigation schedule was developed by [25] in the sugarcane field. Amid sugarcane, developing season Landsat 8 satellite images were gained. The outcome demonstrates that trim water pressure index can be utilized for checking the water pressure and water system planning for sugarcane fields utilizing satellite images with no requirement for ground auxiliary information. For estimating soil moisture SAR derived remote sensing products (Sentinel-1, SRTM and Landsat 8) can be valuable input features [26].

Table 1: Remote sensing summary for estimation of moisture content in soil

Domain	Properties	Pros	Cons
Microwave	Brightness temperature Temperature of soil	Less atmospheric noise Well understood physical	Spatial resolution low Disturbed by surface roughness and vegetation
Optical	Reflection of Soil	Fine spatial resolution coverage	Limited surface penetration Cloud defilement
Thermal IR	Surface temperature	Fine spatial resolution Large coverage Physical very much understood	Constrained surface penetration Cloud defilement Disturbed by meteorological conditions and vegetation

B. Crop water stress monitoring

Identification of crop water stress and investigation of its impact on crop production is essential to advance to decide quantitative loss in crop yield [27-29]. Satellite remote sensing by virtue of its wide coverage, receptivity, and multispectral information becomes an important source for many agricultural applications such as crop condition, crop yield, etc. Momentary variations in status of water within a vineyard are mapped using UAV-based thermal imagery in [30]. The thermal images data turned out to be useful at an occasional time too; in spite of the fact that it did not coordinate regular patterns in status of water yet emulated other process happening at the time of ripening. The reasonableness of a crop water stress index (CWSI) got from very good quality aerial thermal imagery for assessing tree-water status fluctuation in Super High Density (SHD) olive plantations was studied in [31]. Another yield- water stress pointer, the canopy temperature standard deviation inside a thermal image [32] was produced to screen status of crop water utilizing the expectation-maximization algorithm.

Improvement and assessment of thermal IR imaging framework for high spatial and temporal resolution trim water stress observing of corn inside a greenhouse [33] was studied. The outcome demonstrates that the crop water stress index determined by remotely estimating canopy temperature utilizing a thermal infrared imaging framework can be utilized as another water system planning technique to measure spatial and transient soil moisture fluctuation. Recently investigates consolidate thermal imaging with soil remote detecting to upgrade water system administration of sugar beet [34]. In this investigation, it is watched that distinctive water system procedures are required in very heterogeneous plots. An irrigated pinto bean case study on spatial and temporal monitoring is done in [35]. The results demonstrated that the green normalized vegetation index (GNDVI), canopy cover (CC), canopy temperature (CT) were able to differentiate crops with full and deficit irrigation methods at each of the growing region. It showed that the remote sensing method in management of rapid crop stress is much more efficient when compared with the ground-based leaf area index (LAI) estimation.

C. Evapotranspiration and drought stress monitoring

Reliable estimation of ET is of great importance for the calculation of irrigation water requirements, assurance of water spending plan particularly where water resources are scarce and fresh water is a limited resource [36], [37]. Accurate ET estimations in agricultural zones can enhance detection of water stress and aides in irrigation scheduling. Factors affecting evapotranspiration is shown in **Table 2**. Connecting thermal imaging to the physiological indicator in papaya under various water condition was considered in [38]. In pots, plants were developed and subjected to various irrigation treatments. Intelligent irrigation monitoring thermal imaging was recommended by [39]. Thermal imaging in conjunction with the eddy covariance system to portray canopy evapotranspiration (ET) from little heterogeneous grassland was studied in [40]. The outcomes demonstrate that mapped transpiration might be more delicate to streamlined conditions or ecological heterogeneity than stomatal conductance.

Table 2: Factors affecting evapotranspiration

Weather parameters	Crop factors	Environmental conditions
<ul style="list-style-type: none"> • Net radiation • Air temperature • Humidity • Wind Speed 	<ul style="list-style-type: none"> • The difference in resistance to transpiration • Crop height • Crop roughness • Crop rooting characteristics 	<ul style="list-style-type: none"> • Soil water content • Plant density • Ground cover

For arranging proper irrigation management estimation of genuine crop evapotranspiration is exceptionally basic and for that precise estimation of crop coefficient is additionally required. In [41] a satellite-based product coefficient approach is utilized under different vegetation composes and atmosphere conditions. The outcomes demonstrate that it can be used not just to have proper water and farming management yet in addition to investigating and foreseeing crop yield profitability and agrarian drought. Utilizing surface temperature, crop evapotranspiration can be computed. In [42] crop evapotranspiration was figured utilizing two source energy balance (TSEB) which utilized surface temperature estimated by infrared thermometers. The outcome demonstrates that the evapotranspiration computed utilizing infrared thermometers abroad center pivots will be helpful to keep up water efficiency in crops.

LST Retrieval methods from Landsat-8 thermal infrared sensor data is shown in [43]. Assessment of thermal remote sensing lists to evaluate crop evapotranspiration coefficients was done in [44]. To decide the stress coefficient four thermal techniques Crop Water Stress Index – CWSI, Canopy Temperature Ratio – Tcratio, Degrees above Non-Stressed – DANS, Degrees Above Canopy Threshold – DACT were utilized. Results demonstrate that stress location techniques that require fewer data sources are more receptive to water stress. A survey of various satellite-based remote sensing methods is presented in [45]–[47]. It surveys different SEBAL, basic theories, observation methods for evaluating ET from areas with satellite-based land surface temperatures and tabulates the drawbacks of those methods

IV. SOIL PROPERTIES MAPPING

Accurate estimation of variation of soil organic matter is required to accurately estimate soil fertility, structure, porosity, etc. This is required to find out the soil quality as it affects crop production. In [48], [49] studies showed that the timely series satellite data can give proper mapping for soil organic matter in low relief farming areas. Land surface temperature influences the amount of water a soil can hold. Clay soil will be having higher water content and their LST will be low whereas a sandy soil will be having less water holding capacity which in turn results in higher LST [50], [51]. The result in these studies shows that remote sensing using thermal imaging will help in mapping soil properties more accurately which will help farmers for precision agriculture.

Table 3: Soil texture, soil moisture and LST relation

Soil texture	Soil moisture	Land surface temperature
• Sand	• low	• moderate
• Silt	• moderate	• high
• Clay	• high	• low

Soil property mapping using thermal remote sensing helps in providing various information such as crop suitability assessment, land capability classification, irrigation monitoring, etc. Thus, a comparison study between thermal remote sensing and land surveying will help in soil property mapping. **Table 3** shows the soil texture, soil moisture and land surface temperature relation [52]. In [53] author shows thermal remote sensing for classification of hotspots, vegetation, and an assessment technique for use in urban planning. There will be a reduction in land surface temperature with enhancement of urban vegetation cover. Results show that air temperature and LST follows similar patterns at coarse spatial resolution.

V. PLANT DISEASE DETECTION

For proper economic growth and production in agriculture, plant disease should be minimized. It is demonstrated by different researches that thermal imaging helps in identifying the pre-symptomatic impact of illness and pathogen on the plant [54], [55]. The physiological condition of the contaminated tissue, for example, photosynthesis, transpiration, and stomatal conductance are altered during plant-pathogen disease [56]. The decline and increment of the contaminated leaves temperature were clarified in [57] while they were surveying scab illness on apple leaves utilizing thermography and found that the maximum temperature difference (MTD) enhanced with scab advancement and was strongly associated to the measure of disease locales. This MTD can be utilized for infection measurement moreover. Thermal imaging is a vital decision for giving data to plant disease, even though just a couple of studies have utilized this innovation to contemplate and analyze illness and pathogen discovery [58]. Different plant diseases detected by thermal sensors are shown in **Table 4**. Studies show that thermal imaging can be used to determine bruises at the early stage, to certain accuracy insect infestation could be determined.

Table 4: Plant disease detected by thermal sensors

Crop	Disease	Reference
• Grape	• Downy mildew	[59]
• Apple	• Apple Scab	[57]
• Cucumber	• Downy mildew	[60]
• Sugar beet	• Cercospora leaf spot	[61]
• Rose	• Powdery mildew	[62]

VI. CROP MATURITY AND YIELD MAPPING

Despite the fact that few programmed techniques are accessible, crop maturity assessment is accomplished by crop analyzation and visual review. These methodologies are both works escalated and restricting perception number made by a single eyewitness. It is realized that until the point that the

plant achieves the most extreme esteem, the plant physiological esteem increases, and this esteem concurs with the phase of maturity so thermography can be utilized to evaluate the yield development. The assurance of the ideal gather date and the adjusted outline of postharvest hardware [63] should be possible utilizing thermal imaging frameworks. Farmers can do crop insurance, planning of harvest and storage requirements if there is an accurate and early estimation of crop yield. The commonly used method for yield estimation is time-series data models, but it is observed that there is a high deviation in the actual yield from the predicted yield. In this situation thermal imaging can be used for the yield prediction as it is based on the objects own heat radiation.

VII. FIELD TILE MAPPING

There can be extensive damage to crops due to flooding during plant growth periods [64], [65]. Due to excess soil, moisture or high water table during growth season can result in reduced crop growth. By using subsurface drainage this condition can be alleviated. Most of the farmers won't be having the records of the layout of the drainage system in their farms which was laid more than 50 years ago. So, they won't be able to get the benefit of subsurface drainage which will help in improved crop production. Thermal remote sensing helps in addressing these issues by assessing temperature differences within a field [66]. In [67] Remote Sensing based image differencing methods and GIS-based decision tree classification (DTC) were tested for subsurface tile drained area. Studies show that image differencing using remote sensing images provided a more precise classification of tile-drained areas when compared to DTC method. UAS (Unmanned Aircraft System) sensors can be used for drainage pipe mapping [68]. UAS thermal infrared sensors mapped 60% of the drainage pipes in dry soils when compared with unmanned aircraft system visible and near-infrared sensors. In [69], [70] remotely sensed data were used for the identification of subsurface drainage system. But the studies have shown that there is limited success in field tile mapping due to the spectral similarity between crop residue and soil.

VIII. AGRICULTURAL PLASTIC WASTE ESTIMATION

Plastic material use in agriculture has got great benefits but it ends up in large amount of plastic waste to be disposed of [71]. Support Vector Machine was used for image classification with Landsat 8 images. In [72], [73] the efficiency of various image analysis methods applied to hyperspectral satellite data is analyzed. It was used to detect olive oil waste disposal on the island of Crete. A survey of agricultural plastic waste generation and consolidation in Europe is presented in [74]. It represents a systematic effort to map waste analysis. If we are not properly estimating agricultural plastic waste it will create a problem for the future generation. But only very few studies have been done in this field.

IX. CROP RESIDUE COVER AND TILLAGE MAPPING

For creating a protective layer on the agricultural field such as erosion prevention, moisture loss reduction, quality of soil, etc crop residue is essential. Conservation tillage was introduced for the benefit of soil and agricultural environments [75]. Mapping of tillage practices in broad scale is still challenging in thermal remote sensing due to variation in the timing of planting and soil preparation, low revisit time of satellite such as ASTER and Landsat, confounding issues due to soil variation, etc. exact evaluation of crop residue is necessary for proper tillage implementation [76]. Many studies show that thermal imaging technique (both satellite and ground-based) could explain the variability in the amount of crop residue when compared with the visible and near-infrared images [77].

Thermal imaging process plays an important role in many agricultural applications [78]–[80]. While some still need research [81]–[83]; which seems to have great potential in the near future. One such example is ice formation in plant tissues detection, which is challenging, as many methods available were difficult and time-consuming. For the study of freezing behavior and ice-nucleation of potatoes and cauliflower [83], [84] thermal cameras were used. It gave information about ice nucleates and how it spreads into potatoes and cauliflower. Infrared video thermography was used [85] to study the freezing of barley which showed that initial spread of freezing was not problematic but the damage was caused by the second freezing event, which was initiated by the first event.

X. CONCLUSION

This survey has briefly analyzed the current situation of the major application of thermal remote sensing. Thermal imaging systems have been developing quick and assuming an imperative part in different fields of agribusiness. Nonetheless, intensive looks into should be created for its potential application in different procedures of agribusiness, for example, agrarian waste identification, pesticide float in aerial spray applications that are not yet much researched. In spite of the fact that it could be utilized as a part of numerous agricultural operations, it has a few downsides when compared with other imaging techniques; good-quality thermal imaging is costly and to build up an all-inclusive procedure for agrarian applications may not be conceivable since thermal conductivity of crops/fruits shift with change in climatic conditions.

REFERENCES

1. K. Tempfli et al., "Principles of Remote Remote Sensing," p. 591, 2009.
2. D. M. Aubrecht, B. R. Helliker, M. L. Goulden, D. A. Roberts, C. J. Still, and A. D. Richardson, "Continuous, long-term, high-frequency thermal imaging of vegetation: Uncertainties and recommended best practices," *Agric. For. Meteorol.*, vol. 228–229, pp. 315–326, Nov. 2016.
3. C. Lillesand, Kiefer, *Remote Sensing and Image Interpretation*, 6th Edition - Instructor Companion Site. .
4. M. A. Reddy, *Textbook of remote sensing and geographical information systems*. BS Publications, 2013.
5. A. O. Onojeghuo, G. A. Blackburn, J. Huang, D. Kindred, and W. Huang, "Applications of satellite 'hyper-sensing' in Chinese agriculture: Challenges and opportunities," *Int. J. Appl. Earth Obs. Geoinf.*, vol. 64, no. June 2017, pp. 62–86, 2018.
6. D. A. Quattrochi and J. C. Luvall, "Thermal infrared remote sensing for analysis of landscape ecological processes: methods and applications,"

- Landsc. Ecol., vol. 14, no. 6, pp. 577–598, 1999.
7. J. Senthilnath, M. Kandukuri, A. Dokania, and K. N. Ramesh, "Application of UAV imaging platform for vegetation analysis based on spectral-spatial methods," *Comput. Electron. Agric.*, vol. 140, pp. 8–24, Aug. 2017.
8. N. E. Young, R. S. Anderson, S. M. Chignell, A. G. Vorster, R. Lawrence, and P. H. Evangelista, "A survival guide to Landsat preprocessing," *Ecology*, vol. 98, no. 4, pp. 920–932, Apr. 2017.
9. "Landsat 8 Data Users Handbook | Landsat Missions." [Online]. Available: <https://landsat.usgs.gov/landsat-8-data-users-handbook>. [Accessed: 21-Feb-2019].
10. A. Martynenko et al., "Thermal imaging of soybean response to drought stress: the effect of *Ascophyllum nodosum* seaweed extract," *Springerplus*, vol. 5, no. 1, p. 1393, Dec. 2016.
11. J. Berni, P. J. Zarco-Tejada, L. Suarez, and E. Fereres, "Thermal and Narrowband Multispectral Remote Sensing for Vegetation Monitoring From an Unmanned Aerial Vehicle," *IEEE Trans. Geosci. Remote Sens.*, vol. 47, no. 3, pp. 722–738, Mar. 2009.
12. L. Wang and J. J. Qu, "Satellite remote sensing applications for surface soil moisture monitoring: A review," *Front. Earth Sci. China*, vol. 3, no. 2, pp. 237–247, Jun. 2009.
13. A. Soliman et al., "Remote Sensing of Soil Moisture in Vineyards Using Airborne and Ground-Based Thermal Inertia Data," *Remote Sens.*, vol. 5, no. 8, pp. 3729–3748, Jul. 2013.
14. T. Carlson, Carlson, and Toby, "An Overview of the 'Triangle Method' for Estimating Surface Evapotranspiration and Soil Moisture from Satellite Imagery," *Sensors*, vol. 7, no. 8, pp. 1612–1629, Aug. 2007.
15. T. V. Ramachandra, C. Commonwealth of Learning (Vancouver, Indian Institute of Science. Centre for Ecological Sciences., and Karnataka Environment Research Foundation., *Soil and groundwater pollution from agricultural activities*. TERI Press, 2009.
16. N. Agam et al., "An insight to the performance of crop water stress index for olive trees," *Agric. Water Manag.*, vol. 118, pp. 79–86, Feb. 2013.
17. S. Shafian, S. Maas, S. Shafian, and S. J. Maas, "Index of Soil Moisture Using Raw Landsat Image Digital Count Data in Texas High Plains," *Remote Sens.*, vol. 7, no. 3, pp. 2352–2372, Feb. 2015.
18. L. Hassan-Esfahani et al., "Assessment of Surface Soil Moisture Using High-Resolution Multi-Spectral Imagery and Artificial Neural Networks," *Remote Sens.*, vol. 7, no. 3, pp. 2627–2646, Mar. 2015.
19. S. V. Lakshmi, J. James, S. Soundariya, T. Vishalini, and P. K. Pandian, "A Comparison of Soil Texture Distribution and Soil Moisture Mapping of Chennai Coast using Landsat ETM+ and IKONOS Data," *Aquat. Procedia*, vol. 4, pp. 1452–1460, Jan. 2015.
20. P. Rahimzadeh-Bajgiran, A. A. Berg, C. Champagne, and K. Omasa, "Estimation of soil moisture using optical/thermal infrared remote sensing in the Canadian Prairies," *ISPRS J. Photogramm. Remote Sens.*, vol. 83, pp. 94–103, Sep. 2013.
21. Y. Kim, C. J. Still, D. A. Roberts, and M. L. Goulden, "Thermal infrared imaging of conifer leaf temperatures: Comparison to thermocouple measurements and assessment of environmental influences," *Agric. For. Meteorol.*, vol. 248, pp. 361–371, Jan. 2018.
22. F. J. Yandún Narváez, J. Salvo del Pedregal, P. A. Prieto, M. Torres-Torriti, and F. A. Auat Chein, "LiDAR and thermal images fusion for ground-based 3D characterisation of fruit trees," *Biosyst. Eng.*, vol. 151, pp. 479–494, Nov. 2016.
23. S. Khanal, J. Fulton, and S. Shearer, "An overview of current and potential applications of thermal remote sensing in precision agriculture," *Comput. Electron. Agric.*, vol. 139, pp. 22–32, Jun. 2017.
24. K. Ivushkin, H. Bartholomeus, A. K. Bregt, and A. Pulatov, "Satellite Thermography for Soil Salinity Assessment of Cropped Areas in Uzbekistan," *L. Degrad. Dev.*, vol. 28, no. 3, pp. 870–877, Apr. 2017.
25. S. Veysi, A. A. Naseri, S. Hamzeh, and H. Bartholomeus, "A satellite based crop water stress index for irrigation scheduling in sugarcane fields," *Agric. Water Manag.*, vol. 189, pp. 70–86, Jul. 2017.
26. A.-K. Holtgrave, M. Förster, F. Greifeneder, C. Notarnicola, and B. Kleinschmit, "Estimation of Soil Moisture in Vegetation-Covered Floodplains with Sentinel-1 SAR Data Using Support Vector Regression," *PFG – J. Photogramm. Remote Sens. Geoinf. Sci.*, vol. 86, no. 2, pp. 85–101, Apr. 2018.
27. V. Alchanatis et al., "Evaluation of different approaches for estimating and mapping crop water status in cotton with thermal imaging," *Precis. Agric.*, vol. 11, no. 1, pp. 27–41, Feb. 2010.

Thermal Imaging Techniques in Agricultural Applications

28. K. Banerjee, P. Krishnan, and N. Mridha, "Application of thermal imaging of wheat crop canopy to estimate leaf area index under different moisture stress conditions," *Biosyst. Eng.*, vol. 166, pp. 13–27, Feb. 2018.
29. S. Elsayed, M. Elhoweity, H. H. Ibrahim, Y. H. Dewir, H. M. Migdadi, and U. Schmidhalter, "Thermal imaging and passive reflectance sensing to estimate the water status and grain yield of wheat under different irrigation regimes," *Agric. Water Manag.*, vol. 189, pp. 98–110, Jul. 2017.
30. L. G. Santesteban, S. F. Di Gennaro, A. Herrero-Langreo, C. Miranda, J. B. Royo, and A. Matese, "High-resolution UAV-based thermal imaging to estimate the instantaneous and seasonal variability of plant water status within a vineyard," *Agric. Water Manag.*, vol. 183, pp. 49–59, Mar. 2017.
31. G. Egea, C. M. Padilla-Díaz, J. Martínez-Guanter, J. E. Fernández, and M. Pérez-Ruiz, "Assessing a crop water stress index derived from aerial thermal imaging and infrared thermometry in super-high density olive orchards," *Agric. Water Manag.*, vol. 187, pp. 210–221, Jun. 2017.
32. M. Han, H. Zhang, K. C. DeJonge, L. H. Comas, and T. J. Trout, "Estimating maize water stress by standard deviation of canopy temperature in thermal imagery," *Agric. Water Manag.*, vol. 177, pp. 400–409, Nov. 2016.
33. D. L. Mangus, A. Sharda, and N. Zhang, "Development and evaluation of thermal infrared imaging system for high spatial and temporal resolution crop water stress monitoring of corn within a greenhouse," *Comput. Electron. Agric.*, vol. 121, pp. 149–159, Feb. 2016.
34. L. Quebrajo, M. Perez-Ruiz, L. Pérez-Urrestarazu, G. Martínez, and G. Egea, "Linking thermal imaging and soil remote sensing to enhance irrigation management of sugar beet," *Biosyst. Eng.*, vol. 165, pp. 77–87, Jan. 2018.
35. J. Zhou, L. R. Khot, R. A. Boydston, P. N. Miklas, and L. Porter, "Low altitude remote sensing technologies for crop stress monitoring: a case study on spatial and temporal monitoring of irrigated pinto bean," *Precis. Agric.*, vol. 19, no. 3, pp. 555–569, Jun. 2018.
36. I. Petrovic, M. Marjanovic, M. Cosic, S. Savic, and G. Cvijanovic, "Infra-red thermography for detecting drought in agricultural crops and scheduling irrigation," *Ekon. Poljopr.*, vol. 63, no. 2, pp. 461–469, Apr. 2016.
37. [37] R. I. Sholihah et al., "Identification of Agricultural Drought Extent Based on Vegetation Health Indices of Landsat Data: Case of Subang and Karawang, Indonesia," *Procedia Environ. Sci.*, vol. 33, pp. 14–20, Jan. 2016.
38. R. S. Lima et al., "Linking thermal imaging to physiological indicators in *Carica papaya* L. under different watering regimes," *Agric. Water Manag.*, vol. 164, pp. 148–157, Jan. 2016.
39. M. Roopeai, P. Rad, and K.-K. R. Choo, "Cloud of Things in Smart Agriculture: Intelligent Irrigation Monitoring by Thermal Imaging," *IEEE Cloud Comput.*, vol. 4, no. 1, pp. 10–15, Jan. 2017.
40. S. Shimoda and T. Oikawa, "Characteristics of canopy evapotranspiration from a small heterogeneous grassland using thermal imaging," *Environ. Exp. Bot.*, vol. 63, no. 1–3, pp. 102–112, May 2008.
41. J. Park, J. Baik, and M. Choi, "Satellite-based crop coefficient and evapotranspiration using surface soil moisture and vegetation indices in Northeast Asia," *CATENA*, vol. 156, pp. 305–314, Sep. 2017.
42. P. D. Colaizzi, S. A. O'Shaughnessy, S. R. Evett, and R. B. Mounce, "Crop evapotranspiration calculation using infrared thermometers aboard center pivots," *Agric. Water Manag.*, vol. 187, pp. 173–189, Jun. 2017.
43. J. C. Jimenez-Munoz, J. A. Sobrino, D. Skokovic, C. Mattar, and J. Cristobal, "Land Surface Temperature Retrieval Methods From Landsat-8 Thermal Infrared Sensor Data," *IEEE Geosci. Remote Sens. Lett.*, vol. 11, no. 10, pp. 1840–1843, Oct. 2014.
44. E. G. Kullberg, K. C. DeJonge, and J. L. Chávez, "Evaluation of thermal remote sensing indices to estimate crop evapotranspiration coefficients," *Agric. Water Manag.*, vol. 179, pp. 64–73, Jan. 2017.
45. Y.-A. Liou, S. Kar, Y.-A. Liou, and S. K. Kar, "Evapotranspiration Estimation with Remote Sensing and Various Surface Energy Balance Algorithms—A Review," *Energies*, vol. 7, no. 5, pp. 2821–2849, Apr. 2014.
46. M. Abrishamkar and A. Ahmadi, "Evapotranspiration Estimation Using Remote Sensing Technology Based on SEBAL Algorithm," *Iran. J. Sci. Technol. Trans. Civ. Eng.*, vol. 41, no. 1, pp. 65–76, Mar. 2017.
47. A. L. Yagci and J. A. Santanello, "Estimating Evapotranspiration From Satellite Using Easily Obtainable Variables: A Case Study Over the Southern Great Plains, USA," *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, vol. 11, no. 1, pp. 12–23, Jan. 2018.
48. M.-S. Zhao, G.-L. Zhang, F. Liu, D.-C. Li, and Y.-G. Zhao, "Mapping Soil Organic Matter in Low-Relief Areas Based on Time Series Land Surface Diurnal Temperature Difference," Springer, Singapore, 2016, pp. 47–62.
49. A. Eisele et al., "Applicability of the Thermal Infrared Spectral Region for the Prediction of Soil Properties Across Semi-Arid Agricultural Landscapes," *Remote Sens.*, vol. 4, no. 11, pp. 3265–3286, Oct. 2012.
50. D.-C. Wang et al., "Retrieval and Mapping of Soil Texture Based on Land Surface Diurnal Temperature Range Data from MODIS," *PLoS One*, vol. 10, no. 6, p. e0129977, Jun. 2015.
51. E. Aldana-Jague, G. Heckrath, A. Macdonald, B. van Wesemael, and K. Van Oost, "UAS-based soil carbon mapping using VIS-NIR (480–1000 nm) multi-spectral imaging: Potential and limitations," *Geoderma*, vol. 275, pp. 55–66, Aug. 2016.
52. W. J. Northcott, A. K. Verma, and R. A. C. Cooke, "Mapping Subsurface Drainage Systems using Remote Sensing and GIS," pp. 2625–2634, 01-Dec-2000.
53. A. M. Coutts, R. J. Harris, T. Phan, S. J. Livesley, N. S. G. Williams, and N. J. Tapper, "Thermal infrared remote sensing of urban heat: Hotspots, vegetation, and an assessment of techniques for use in urban planning," *Remote Sens. Environ.*, vol. 186, pp. 637–651, Dec. 2016.
54. A.-K. Mahlein, E.-C. Oerke, U. Steiner, and H.-W. Dehne, "Recent advances in sensing plant diseases for precision crop protection," *Eur. J. Plant Pathol.*, vol. 133, no. 1, pp. 197–209, May 2012.
55. A.-K. Mahlein, "Plant Disease Detection by Imaging Sensors – Parallels and Specific Demands for Precision Agriculture and Plant Phenotyping," *Plant Dis.*, vol. 100, no. 2, pp. 241–251, Feb. 2016.
56. H. Xu, S. Zhu, Y. Ying, and H. Jiang, "Early detection of plant disease using infrared thermal imaging," 2006, vol. 6381, p. 638110.
57. E.-C. Oerke, P. Fröhling, and U. Steiner, "Thermographic assessment of scab disease on apple leaves," *Precis. Agric.*, vol. 12, no. 5, pp. 699–715, Oct. 2011.
58. R. Ishimwe, K. Abutaleb, and F. Ahmed, "Applications of Thermal Imaging in Agriculture—A Review," *Adv. Remote Sens.*, vol. 03, no. 03, pp. 128–140, 2014.
59. M. Stoll, H. R. Schultz, G. Baecker, and B. Berkelmann-Loehnertz, "Early pathogen detection under different water status and the assessment of spray application in vineyards through the use of thermal imagery," *Precis. Agric.*, vol. 9, no. 6, pp. 407–417, Dec. 2008.
60. E.-C. Oerke, U. Steiner, H.-W. Dehne, and M. Lindenthal, "Thermal imaging of cucumber leaves affected by downy mildew and environmental conditions," *J. Exp. Bot.*, vol. 57, no. 9, pp. 2121–2132, Jan. 2006.
61. L. Chaerle, D. Hagenbeek, E. De Bruyne, R. Valcke, and D. Van Der Straeten, "Thermal and Chlorophyll-Fluorescence Imaging Distinguish Plant-Pathogen Interactions at an Early Stage," *Plant Cell Physiol.*, vol. 45, no. 7, pp. 887–896, Jul. 2004.
62. M. Jafari, S. Minaei, N. Safaei, and F. Torkamani-Azar, "Early detection and classification of powdery mildew-infected rose leaves using ANFIS based on extracted features of thermal images," *Infrared Phys. Technol.*, vol. 76, pp. 338–345, May 2016.
63. M. Linke, H. Beuche, M. Geyer, and H. J. Hellebrand, "Possibilities and limits of the use of thermography for the examination of horticultural products," vol. 6, no. 2. Landwirtschaftsverl., 2000.
64. K. W. King, M. R. Williams, and N. R. Fausey, "Contributions of Systematic Tile Drainage to Watershed-Scale Phosphorus Transport," *J. Environ. Qual.*, vol. 44, no. 2, p. 486, Mar. 2015.
65. D. R. Smith et al., "Surface Runoff and Tile Drainage Transport of Phosphorus in the Midwestern United States," *J. Environ. Qual.*, vol. 44, no. 2, p. 495, Mar. 2015.
66. X. Song, J. Wang, W. Huang, L. Liu, G. Yan, and R. Pu, "The delineation of agricultural management zones with high resolution remotely sensed data," *Precis. Agric.*, vol. 10, no. 6, pp. 471–487, Dec. 2009.
67. K. Gökkaya, M. Budhathoki, S. F. Christopher, B. R. Hanrahan, and J. L. Tank, "Subsurface tile drained area detection using GIS and remote sensing in an agricultural watershed," *Ecol. Eng.*, vol. 108, pp. 370–379, Nov. 2017.
68. B. Allred, N. Eash, R. Freeland, L. Martinez, and D. B. Wishart, "Effective and efficient agricultural drainage pipe mapping with UAS thermal infrared imagery: A case study," *Agric. Water Manag.*, vol. 197, pp. 132–137, 2018.
69. B. S. B. S. Naz and L. C. L. C. Bowling, "Automated Identification of Tile Lines from Remotely Sensed Data," *Trans. ASABE*, vol. 51, no. 6, pp. 1937–1950, 2008.

70. B. S. Naz, S. Ale, and L. C. Bowling, "Detecting subsurface drainage systems and estimating drain spacing in intensively managed agricultural landscapes," *Agric. Water Manag.*, vol. 96, no. 4, pp. 627–637, Apr. 2009.
71. A. Lanorte et al., "Agricultural plastic waste spatial estimation by Landsat 8 satellite images," *Comput. Electron. Agric.*, vol. 141, pp. 35–45, Sep. 2017.
72. [72] A. Agapiou, N. Papadopoulos, and A. Sarris, "Monitoring olive mills waste disposal areas in Crete using very high resolution satellite data," *Egypt. J. Remote Sens. Sp. Sci.*, vol. 19, no. 2, pp. 285–295, Dec. 2016.
73. D. D. Alexakis, A. Sarris, C. Kalaitzidis, N. Papadopoulos, and P. Soupios, "Integrated use of satellite remote sensing, GIS, and ground spectroscopy techniques for monitoring olive oil mill waste disposal areas on the island of Crete, Greece," *Int. J. Remote Sens.*, vol. 37, no. 3, pp. 669–693, Feb. 2016.
74. D. Briassoulis et al., "Review, mapping and analysis of the agricultural plastic waste generation and consolidation in Europe," *Waste Manag. Res.*, vol. 31, no. 12, pp. 1262–1278, Dec. 2013.
75. B. Zheng, J. B. Campbell, G. Serbin, and J. M. Galbraith, "Remote sensing of crop residue and tillage practices: Present capabilities and future prospects," *Soil Tillage Res.*, vol. 138, pp. 26–34, May 2014.
76. "Mapping Crop Residue and Tillage Intensity on Chesapeake Bay Farmland | Land Imaging Report Site." [Online]. Available: <https://eros.usgs.gov/doi-remote-sensing-activities/2015/usgs/mapping-crop-residue-and-tillage-intensity-chesapeake-bay-farmland>. [Accessed: 06-Mar-2019].
77. B. Zheng, J. B. Campbell, G. Serbin, and C. S. T. Daughtry, "Multitemporal remote sensing of crop residue cover and tillage practices: A validation of the minNDTI strategy in the United States," *J. Soil Water Conserv.*, vol. 68, no. 2, pp. 120–131, Mar. 2013.
78. M. Bisquert, J. M. Sánchez, R. López-Urrea, and V. Caselles, "Estimating high resolution evapotranspiration from disaggregated thermal images," *Remote Sens. Environ.*, vol. 187, pp. 423–433, Dec. 2016.
79. R. A. ElGamal, S. S. Kishk, and G. M. ElMasry, "Validation of CFD models for the deep-bed drying of rice using thermal imaging," *Biosyst. Eng.*, vol. 161, pp. 135–144, Sep. 2017.
80. S. M. Z. Younis and J. Iqbal, "Estimation of soil moisture using multispectral and FTIR techniques," *Egypt. J. Remote Sens. Sp. Sci.*, vol. 18, no. 2, pp. 151–161, Dec. 2015.
81. A. Nishar, S. Richards, D. Breen, J. Robertson, and B. Breen, "Thermal infrared imaging of geothermal environments and by an unmanned aerial vehicle (UAV): A case study of the Wairakei – Tauhara geothermal field, Taupo, New Zealand," *Renew. Energy*, vol. 86, pp. 1256–1264, Feb. 2016.
82. L. Jiao, D. Dong, H. Feng, X. Zhao, and L. Chen, "Monitoring spray drift in aerial spray application based on infrared thermal imaging technology," *Comput. Electron. Agric.*, vol. 121, pp. 135–140, Feb. 2016.
83. M. J. Fuller and M. Wisniewski, "The use of infrared thermal imaging in the study of ice nucleation and freezing of plants," *J. Therm. Biol.*, vol. 23, no. 2, pp. 81–89, Apr. 1998.
84. L. Cuiibus, M. Castro-Giráldez, P. J. Fito, and A. Fabbri, "Application of infrared thermography and dielectric spectroscopy for controlling freezing process of raw potato," *Innov. Food Sci. Emerg. Technol.*, vol. 24, pp. 80–87, Aug. 2014.
85. R. S. Pearce and M. P. Fuller, "Freezing of barley studied by infrared video thermography," *Plant Physiol.*, vol. 125, no. 1, pp. 227–40, Jan. 2001.

has published 55 papers in peer reviewed national and international journals, 62 papers in reputed national and international conferences and 3 book chapters. Citations and h-index (As on July 2019): Google Scholar, 844 citations (h-index = 15 and i-10index = 22), Scopus – 564 (h-index = 14), Web of Science – 369 (h-index=11). He is involved in research which covers various sorts of wireless networks such as ad hoc networks, sensor networks, mesh networks, network security, smart antennas, Index modulation, etc. He has guided 5 PhD's and currently, he is guiding 6 PhD students. He is a reviewer of various reputed journals and conferences and chaired many conferences. He is a Fellow IETE (FIETE) India, member IE (MIE) India, member ISTE (MISTE) India and senior member of IEEE (SMIEEE), USA and member of IACSIT. He has been listed in Marquis' Who's Who in the World (2012 Edition), USA and Top 100 Engineers by IBC, UK.



Mohammed Ahamed J is a Scientist at RRSC-S, National Remote Sensing Centre, Indian Space Research Organization. He has done his M.Tech in Satellite Remote Sensing and GIS from Indian Institute of Remote Sensing, M.Sc and B.Sc from University of Agricultural Science GKVK, Bangalore. Currently he is working on Satellite Remote Sensing applications in the field of natural resource management (Agriculture crop growth and monitoring, Drought Assessment and Monitoring, Horticulture applications and Agro-meteorology)

AUTHORS PROFILE



Aryalekshmi B N has 3 years of teaching and research experience. She has experience as project trainee in ISRO (RRSC), DRDO and TataElxsi. She has publications in reputed national/international journals and conferences. She is a member of IEEE and IEEE Geoscience and Remote Sensing Society. She is a Lifetime member in IETE. Currently, she is pursuing her Ph.D.



Dr. Rajashekhar C. Biradar has 29 years of teaching and research experience. He has many research publications in reputed national/international journals and conferences. Some of the journals where his research articles published are Elsevier, IET, Springer, Wiley and IOS Press, having good impact factors. He